

Measuring $\psi'' \rightarrow K_S^0 K_L^0$ as a test of the S - and D -wave mixing of charmonia

P. Wang,* X. H. Mo, and C. Z. Yuan
Institute of High Energy Physics, CAS, Beijing 100039, China
 (Dated: March 13, 2017)

Adding to the long standing “ $\rho\pi$ puzzle” in ψ' and J/ψ decays, recently BEIJING Spectrometer (BES) reported $\mathcal{B}(\psi' \rightarrow K_S^0 K_L^0)$ which is enhanced relative to the pQCD “12% rule” expectation from $\mathcal{B}(J/\psi \rightarrow K_S^0 K_L^0)$. If the enhancement is due to the mixing of the S - and D -wave charmonium states as in the $\rho\pi$ case, the newly measured $\mathcal{B}(\psi' \rightarrow K_S^0 K_L^0)$ gives a constraint on $\mathcal{B}(\psi'' \rightarrow K_S^0 K_L^0)$. It serves as a good test for the scenario of the S - and D -wave mixing in the ψ' and ψ'' .

PACS numbers: 12.39.Pn, 13.25.Gv, 14.40.Gx

I. INTRODUCTION

From the perturbative QCD (pQCD), it is expected that both J/ψ and $\psi(3686)$ (shortened as ψ') decaying into light hadrons are dominated by the annihilation of $c\bar{c}$ into three gluons, with widths proportional to the square of the wave function at the origin [1]. This yields the pQCD “12% rule”, that is

$$Q_h = \frac{\mathcal{B}_{\psi' \rightarrow h}}{\mathcal{B}_{J/\psi \rightarrow h}} = \frac{\mathcal{B}_{\psi' \rightarrow e^+ e^-}}{\mathcal{B}_{J/\psi \rightarrow e^+ e^-}} \approx 12\% . \quad (1)$$

The violation of the above rule was first observed in $\rho\pi$ and $K^{*+}K^- + c.c.$ modes by Mark II [2], since then BES has measured many two-body decay modes of ψ' , among which some obey the 12% rule while others violate it [3]. There have been many theoretical efforts trying to solve the puzzle [4], however, none explains all the existing experimental data satisfactorily and naturally [5].

A most recent explanation of the “ $\rho\pi$ puzzle” using the S - and D -wave charmonia mixing was proposed by Rosner [6]. In this scheme, the mixing of $\psi(2^3S_1)$ state and $\psi(1^3D_1)$ is in such a way which leads to almost complete cancellation of the decay amplitude of $\psi' \rightarrow \rho\pi$, and the missing $\rho\pi$ decay mode of ψ' shows up instead as enhanced decay mode of $\psi(3770)$ (shortened as ψ''). A study on the measurement of $\psi'' \rightarrow \rho\pi$ in e^+e^- experiments shows that with the decay rate predicted by the S - and D -wave mixing, the destructive interference between the three-gluon decay amplitude of the ψ'' resonance and the continuum one-photon amplitude leads to a very small cross section [7], which is in agreement with the unpublished upper limit of the $\rho\pi$ cross section at the ψ'' peak by Mark III [8]. Although this needs to be further tested by high luminosity experiment operating at the ψ'' mass energy, such as CLEO-c, it already showed that $\mathcal{B}(\psi'' \rightarrow \rho\pi)$ is most probably at the order of 10^{-4} , in agreement with the prediction of the S - and D -wave mixing scheme.

If the S - and D -wave mixing is the key for solving the $\rho\pi$ puzzle, it applies to other decay modes as well, such

as pseudoscalar pseudoscalar (PP) mode like $K_S^0 K_L^0$. Recently, BES collaboration reported the $K_S^0 K_L^0$ branching ratios of J/ψ and ψ' decays [9, 10]:

$$\begin{aligned} \mathcal{B}(J/\psi \rightarrow K_S^0 K_L^0) &= (1.82 \pm 0.04 \pm 0.13) \times 10^{-4} , \\ \mathcal{B}(\psi' \rightarrow K_S^0 K_L^0) &= (5.24 \pm 0.47 \pm 0.48) \times 10^{-5} . \end{aligned} \quad (2)$$

These results yield $Q_{K_S^0 K_L^0} = (28.8 \pm 3.7)\%$, which is enhanced relative to the 12% rule by more than 4σ . In this paper, the $\psi' \rightarrow K_S^0 K_L^0$ enhancement is explained in the S - and D -wave charmonia mixing scheme, and the $\psi'' \rightarrow K_S^0 K_L^0$ decay rate is estimated with the inputs $\mathcal{B}(J/\psi \rightarrow K_S^0 K_L^0)$ and $\mathcal{B}(\psi' \rightarrow K_S^0 K_L^0)$. In following sections, the mixing scheme is introduced briefly, then the branching ratio of $\psi'' \rightarrow K_S^0 K_L^0$ is calculated with the measured e^+e^- and $K_S^0 K_L^0$ decay rates of J/ψ and ψ' , assuming the mixing of S - and D -wave. Finally the experiment search for $\psi'' \rightarrow K_S^0 K_L^0$ is proposed.

II. S - AND D -WAVE MIXING SCHEME

To explain the measured Γ_{ee} of ψ'' , it is suggested [11–13] that the mass eigenstates ψ' and ψ'' are the mixtures of the S - and D -wave of charmonia, namely $\psi(2^3S_1)$ and $\psi(1^3D_1)$ states. In this scheme,

$$\begin{aligned} |\psi'\rangle &= |2^3S_1\rangle \cos\theta - |1^3D_1\rangle \sin\theta , \\ |\psi''\rangle &= |2^3S_1\rangle \sin\theta + |1^3D_1\rangle \cos\theta , \end{aligned} \quad (3)$$

where θ is the mixing angle between pure $\psi(2^3S_1)$ and $\psi(1^3D_1)$ states and is fitted from the leptonic widths of ψ'' and ψ' to be either $(-27 \pm 2)^\circ$ or $(12 \pm 2)^\circ$ [6]. The latter value of θ is consistent with the coupled channel estimates [11, 14] and with the ratio of ψ' and ψ'' partial widths to $J/\psi\pi^+\pi^-$ [12]. Hereafter, the discussions in this paper are solely for the mixing angle $\theta = 12^\circ$.

As in the discussion of Ref. [6], since both hadronic and leptonic decay rates are proportional to the square of the wave function at the origin $|\Psi(0)|^2$, it is expected that if ψ' is a pure $\psi(2^3S_1)$ state, then for any hadronic final states f ,

$$\Gamma(\psi' \rightarrow f) = \Gamma(J/\psi \rightarrow f) \frac{\Gamma(\psi' \rightarrow e^+e^-)}{\Gamma(J/\psi \rightarrow e^+e^-)} . \quad (4)$$

*Electronic address: wangp@mail.ihep.ac.cn

The electronic partial width of J/ψ is expressed in potential model by [15]

$$\Gamma(J/\psi \rightarrow e^+e^-) = \frac{4\alpha^2 e_c^2}{M_{J/\psi}^2} |R_{1S}(0)|^2, \quad (5)$$

with α the QED fine structure constant, $e_c = 2/3$, $M_{J/\psi}$ the J/ψ mass and $R_{1S}(0)$ the radial 1^3S_1 wave function at the origin.

ψ' is not a pure $\psi(2^3S_1)$ state, its electronic partial width is expressed as [6]

$$\begin{aligned} \Gamma(\psi' \rightarrow e^+e^-) &= \frac{4\alpha^2 e_c^2}{M_{\psi'}^2} \\ &\times \left| \cos \theta R_{2S}(0) - \frac{5}{2\sqrt{2}m_c^2} \sin \theta R_{1D}''(0) \right|^2, \end{aligned} \quad (6)$$

with $M_{\psi'}$ the ψ' mass, $R_{2S}(0)$ the radial 2^3S_1 wave function at the origin and $R_{1D}''(0)$ the second derivative of the radial 1^3D_1 wave function at the origin.

If Eq. (4) holds for a pure 2^3S_1 state, $\psi'' \rightarrow f$, $\psi' \rightarrow f$ and $J/\psi \rightarrow f$ partial widths are to be

$$\begin{aligned} \Gamma(\psi'' \rightarrow f) &= \frac{C_f}{M_{\psi''}^2} |\sin \theta R_{2S}(0) + \eta \cos \theta|^2, \\ \Gamma(\psi' \rightarrow f) &= \frac{C_f}{M_{\psi'}^2} |\cos \theta R_{2S}(0) - \eta \sin \theta|^2, \\ \Gamma(J/\psi \rightarrow f) &= \frac{C_f}{M_{J/\psi}^2} |R_{1S}(0)|^2, \end{aligned} \quad (7)$$

where C_f is a common factor for the final state f , $M_{\psi''}$ the ψ'' mass, and $\eta = |\eta|e^{i\phi}$ is a complex parameter with ϕ being the relative phase between $\langle f | 1^3D_1 \rangle$ and $\langle f | 2^3S_1 \rangle$.

III. UPPER AND LOWER BOUNDS OF $\mathcal{B}(\psi'' \rightarrow K_S^0 K_L^0)$

With Eqs. (5, 6, 7), the following two equations are derived :

$$\begin{aligned} \frac{\Gamma(\psi' \rightarrow f)}{\Gamma(J/\psi \rightarrow f)} &= \frac{\Gamma(\psi' \rightarrow e^+e^-)}{\Gamma(J/\psi \rightarrow e^+e^-)} \\ &\times \left| \frac{\cos \theta R_{2S}(0) - \eta \sin \theta}{\cos \theta R_{2S}(0) - \frac{5}{2\sqrt{2}m_c^2} \sin \theta R_{1D}''(0)} \right|^2, \end{aligned} \quad (8)$$

and

$$\frac{\Gamma(\psi'' \rightarrow f)}{\Gamma(\psi' \rightarrow f)} = \frac{M_{\psi'}^2}{M_{\psi''}^2} \left| \frac{\sin \theta R_{2S}(0) + \eta \cos \theta}{\cos \theta R_{2S}(0) - \eta \sin \theta} \right|^2. \quad (9)$$

It is easy to see that if $\theta = 0$, i.e. ψ' were a pure $\psi(2^3S_1)$ state, Eq. (8) becomes Eq. (4).

In the following, the discussion focuses on $f = K_S^0 K_L^0$ final state. The partial widths of ψ' and J/ψ to e^+e^- [16] and $K_S^0 K_L^0$ [9, 10] are all measured by experiments; $R_{2S}(0) = 0.734 \text{ GeV}^{3/2}$ and $5R_{1D}''(0)/(2\sqrt{2}m_c^2) = 0.095 \text{ GeV}^{3/2}$ are given in Ref. [6], so for the final state $K_S^0 K_L^0$, Eq. (8) has only one unknown variable η . Since η is complex, for any given phase, its module can be determined. Then with η substituting into Eq. (9), $\Gamma(\psi'' \rightarrow K_S^0 K_L^0)$ can be calculated.




FIG. 1: The variation of $\mathcal{B}(\psi'' \rightarrow K_S^0 K_L^0) \times 10^5$ against the phase ϕ (in degree).

Since the phase of η is a free parameter, so the decay rate of $\psi'' \rightarrow K_S^0 K_L^0$ is constrained in a range. According to Eqs. (8) and (9), the variation of branching ratio against the phase is shown in Fig. 1, from which we see that

$$0.12 \pm 0.07 \leq 10^5 \times \mathcal{B}(\psi'' \rightarrow K_S^0 K_L^0) \leq 3.8 \pm 1.1. \quad (10)$$

Here the upper bound corresponds to $\phi = 0^\circ$ and the lower bound to $\phi = \pm 180^\circ$. The uncertainties are due to the mixing angle θ , the measurements of $\mathcal{B}(\psi' \rightarrow K_S^0 K_L^0)$ and $\mathcal{B}(J/\psi \rightarrow K_S^0 K_L^0)$, with the first two dominate.

IV. EXPERIMENTAL TEST

It is instructive to look at the range of the phase ϕ from other decay modes, such as $\rho\pi$. The recent phenomenological estimation [17] gives the branching ratio of $\psi' \rightarrow \rho\pi$ at the level of 10^{-4} , which indicates the almost complete cancellation between $\cos \theta R_{2S}(0)$ and $\eta \sin \theta$ in Eq. (7). In another word, the small $\mathcal{B}(\psi' \rightarrow \rho\pi)$ means the phase ϕ of η is around zero. With incomplete cancellation between $\cos \theta R_{2S}(0)$ and $\eta \sin \theta$ which results in $\mathcal{B}(\psi' \rightarrow \rho\pi) = 1.11 \times 10^{-4}$ [17], and latest results by BES of $\mathcal{B}(J/\psi \rightarrow \rho\pi) \sim 2.1\%$ [18], ϕ is constrained to be less than 11° . As a pedagogical guess, ϕ is expected to be small for other decay modes too. In such case, the prediction $\mathcal{B}(\psi'' \rightarrow K_S^0 K_L^0)$ would be close to the upper bound in Eq. (10), that is

$$\mathcal{B}(\psi'' \rightarrow K_S^0 K_L^0) \approx (3.8 \pm 1.1) \times 10^{-5}. \quad (11)$$

Currently, BES has accumulated about 20 pb^{-1} data while CLEO-c has collected 55 pb^{-1} data at ψ'' peak. By virtue of Eq. (11), assuming 40% efficiency for detecting $K_S^0 \rightarrow \pi^+ \pi^-$, then one expects 1.7 events from BES and 4.6 events from CLEO-c. Utilizing these samples, most probably an upper limit can be set by BES, while the signal can be seen at CLEO-c. With the expected larger ψ'' data sample of several fb^{-1} [19] in immediate future, CLEO-c can give a definite answer for prediction of Eq. (11), or test the lower bound of Eq. (10) in case the phase ϕ is not small.

V. DISCUSSION

In the S - and D -wave mixing scheme, the observed $\psi' \rightarrow K_S^0 K_L^0$ enhancement relative to the 12 % rule implies a $\psi'' \rightarrow K_S^0 K_L^0$ decay branching ratio at the order of 10^{-6} to 10^{-5} . So the measurement of $\mathcal{B}(\psi'' \rightarrow K_S^0 K_L^0)$ will provide a clear-cut test of the S - and D -wave mixing scenario.

Unlike the $\rho\pi$ modes, $K_S^0 K_L^0$ mode of the 1^{--} charmonium decay is only through strong interaction due to SU(3) symmetry [20]. There is no complication of electromagnetic interaction and continuum one-photon annihilation as well as the interference between them [21]. So the observed $K_S^0 K_L^0$ in e^+e^- experiment is completely from resonance decays.

If the ψ' and ψ'' are indeed the S - and D -wave charmonia mixtures, not only the vector pseudoscalar [6] and the

pseudoscalar pseudoscalar modes will be affected, but all the other modes in ψ' decays will be affected as well, such as vector tensor, axial-vector pseudoscalar and so forth. For the decay modes which have been measured both at ψ' and J/ψ , the corresponding branching ratio at ψ'' can be evaluated under the assumption of pQCD. Then the measurements at ψ'' provide a test for the mixing scheme, at the same time help to reveal the charmonium decay dynamics and the relation between J/ψ and ψ' decays.

The mixing scheme is a simple and natural model, it will provide a new angle of purview of understanding the $\rho\pi$ puzzle between J/ψ and ψ' decays, and the non- $D\bar{D}$ decay of ψ'' .

VI. SUMMARY

In this paper, the S - and D -wave mixing scheme of charmonium states is applied on $\psi' \rightarrow K_S^0 K_L^0$ to explain its enhancement relative to the pQCD 12% rule, and the branching ratio of $\psi'' \rightarrow K_S^0 K_L^0$ is predicted. It is suggested that with the data samples collected currently and the larger data sample expected from CLEO-c soon, the mixing scheme is to be tested.

Acknowledgments

This work is supported by 100 Talents Program of the Chinese Academy of Sciences under contract No. U-25.

-
- [1] T. Appelquist and H. D. Politzer, Phys. Rev. Lett. **34**, 43 (1975); A. De Rújula and S. L. Glashow, Phys. Rev. Lett. **34**, 46 (1975).
 - [2] M. E. B. Franklin *et al.*, Phys. Rev. Lett. **51**, 963 (1983).
 - [3] For a recent review, see Gu Yifan, "Experimental Study on Charmonium Physics at Beijing Electro-Positron Collider" in "China Basic Science(2002.2)" (in Chinese), page 7-17.
 - [4] W. S. Hou and A. Soni, Phys. Rev. Lett. **50**, 569 (1983); S. J. Brodsky and M. Karliner, Phys. Rev. Lett. **78**, 4682 (1997); M. Chaichian and N. A. Törnqvist, Nucl. Phys. B **323**, 75 (1989); S. S. Pinsky, Phys. Lett. B **236**, 479 (1990); G. Karl and W. Roberts, Phys. Lett. B **144**, 243 (1984); X. Q. Li, D. V. Bugg and B. S. Zou, Phys. Rev. D **55**, 1421 (1997); J. M. Gérard and J. Weyers, Phys. Lett. B **462**, 324 (1999); T. Feldmann and P. Kroll, Phys. Rev. D **62**, 074006 (2000).
 - [5] For recent reviews, see for example M. Suzuki, Phys. Rev. D **63**, 054021 (2001); Y. F. Gu and X. H. Li, Phys. Rev. D **63**, 114019 (2001).
 - [6] J. L. Rosner, Phys. Rev. D **64**, 094002 (2001).
 - [7] P. Wang, X. H. Mo and C. Z. Yuan, Phys. Lett. B **574**, 41 (2003).
 - [8] Yanong Zhu, Ph. D. thesis, California Institute of Technology, 1988, Caltech Report No. CALT-68-1513.
 - [9] BES Collaboration, J. Z. Bai *et al.*, Phys. Rev. D **69**, 012003 (2004).
 - [10] BES Collaboration, J. Z. Bai *et al.*, Phys. Rev. Lett. **92**, 052001 (2004).
 - [11] E. Eichten, K. Gottfried, T. Kinoshita, K. D. Lane and T.-M. Yan, Phys. Rev. D **17**, 3090 (1978); D **21**, 313(E) (1980); D **21**, 203 (1980).
 - [12] Y.-P. Kuang and T. M. Yan, Phys. Rev. D **41**, 155 (1990).
 - [13] J. L. Rosner, in Particles and Fields 3, Proceedings of Banff Summer Institute (CAP) 1988, Banff, Alberta, 14-26 August 1988, edited by A. N. Kamal and F. C. Khanna (World Scientific, Singapore, 1989), p. 395.
 - [14] K. Heikkilä, N. A. Törnqvist and S. Ono, Phys. Rev. D **29**, 110 (1984); D **29**, 2136(E) (1984).
 - [15] V. A. Novikov *et al.*, Phys. Rep. **41**, 1 (1978).
 - [16] Particle Data Group, K. Hagiwara *et al.*, Phys. Rev. D **66**, 010001 (2002).
 - [17] P. Wang, C. Z. Yuan and X. H. Mo, Phys. Rev. D **69**, 057502 (2004).
 - [18] BES Collaboration, J. Z. Bai *et al.*, hep-ex/0402013; The $\rho\pi$ branching ratio could be different from $\pi^+ \pi^- \pi^0$ branching ratio, due to the interference with the excited ρ states as shown in reference SLAC-PUB-5674 (1991) by L. P. Chen and W. Dunwoodie.
 - [19] R. A. Briere, Recent results from CLEO, talk at the 8th international workshop on meson production, properties and interactions, Cracow, Poland, June 4-8, 2004, <http://www-hep.phys.cmu.edu/users/rbriere/Meson2004.ppt>.

- [20] H. E. Haber and J. Perrier, Phys. Rev. D **32**, 2961 (1985).
- [21] P. Wang, C. Z. Yuan, X. H. Mo and D. H. Zhang, Phys.

Lett. B **593**, 89 (2004).